



Growth Equations in Forest Research: Mathematical Basis and Model Similarities

Christian Salas-Eljatib^{1,2,3} · Lauri Mehtätalo⁴ · Timothy G. Gregoire⁵ · Daniel P. Soto⁶ · Rodrigo Vargas-Gaete^{7,8}

Accepted: 24 August 2021

© The Author(s), under exclusive licence to Springer Nature Switzerland AG 2021

Abstract

Purpose of Review Growth equations have been widely used in forest research, commonly to assess ecosystem-level behavior and forest management. Nevertheless, the large number of growth equations has obscured the growth-rate behavior of each of these equations and several different terms for referring to common phenomena. This review presents a unified mathematical treatment of growth-rates besides several well-known growth equations by giving their mathematical basis and representing their behavior using tree growth data as an example.

Recent Findings We highlight the mathematical differences among several growth equations that can be better understood by using their differential equations forms rather than their integrated forms. Moreover, the assumed-and-claimed biological basis of these growth-rate models has been taken too seriously in forest research. The focus should be on using a plausible equation for the organism being modelled. We point out that more attention should be drawn to parameter estimation strategies and behavior analysis of the proposed models. Thus, it is difficult for a single model to capture all possible shapes and rates that such a complex biological process as tree growth can depict in nature.

Summary We pointed out misleading concepts attributed to some growth equations; however, the differences come from their mathematical properties rather than pure biological reasoning. Using the tree growth data, we depict those differences. Thus, comparisons of some functional forms (at least simple ones) must be carried out before selecting a function for drawing scientific findings.

Keywords Differential equations · Growth-rates · von Bertalanffy · Logistic · Trees

Introduction

Growth is a term which everybody understands but not necessarily in the same way. Although we can track definitions of growth as far as Aristotle [1], one of the first obstacles in understanding tree growth is the lack of precision in the definition of what is meant by growth. Growth is determined by cell division, cell extension and cell differentiation [2].

Ergo, growth is a phenomenon that captures the interest of research in a suite of disciplines [3], especially in plant sciences [4]. The growth pattern in forest trees is divided into primary, i.e., the growth from a bud, root, tip or another apical meristem, and secondary, i.e., growth from the cambium [5]. Growth implies an increase in size and the formation of new tissue; however, it growth may occur when older organs, particularly leaves, are dropping off faster than new ones are being formed [6]. The term *net growth* includes also this component, whereas *gross growth* does not. In forestry, these terms are mainly used at the stand level, and using volume as variable of reference, as follows: net growth refers to the difference between volume at the end of the period and at the beginning; gross growth must also incorporate the growth of trees which may have died in the period [5, 7]. We refer here to the growth of variables such as height and stem diameters as the irreversible increase in

This article is part of the Topical Collection on *Modelling Productivity and Function*

✉ Christian Salas-Eljatib
cseljatib@gmail.com

Extended author information available on the last page of the article.

leading-shoot length and stem diameter, respectively [8], or in a more physiological context, the incorporation of carbon into structural material [2]. Nevertheless, we follow [9], in the sense that the level of abstraction of our modelling framework emphasizes the physically measurable exterior tree characteristics, which we call *variables*.

Let us take a closer look at the use of the terms “growth”, “increment” and “yield” in forestry. Bruce [10] stated, using height growth equations as an example, that there are *growth* and *growth-rate* equations, the first one being an expression giving the variable as a function of time, and the second as the differential of the growth equation. He also emphasized that growth-rate is determined at one point in time. In forestry, growth over a time period shorter than a year may be confusing because the growth usually occurs during summertime. Yang et al. [11] defined the term *increment* as the increase in size of an organism within a certain time interval, i.e., as the difference in observed or predicted growth at two points in time [10]. *Current annual increment (CAI)* is the difference in the growth at the beginning and the end of the year and *periodic annual increment (PAI)* for longer time units [12]. Term *yield* has also been used in the same meaning as growth [12–16], but it has mainly been associated with volume. For example, Clutter [14] and Curtis [15] refer to growth-rate as the derivative form and to yield as the integral of volume. Clutter [14] defined that if integrating the growth-rate equation gives the yield equation, the models are compatible. Even though the forestry literature often recognizes the works by [12] and [14] as the first to mention the consistency between growth-rate equations and the yield equations, many researchers in plant ecology [17–19], animal growth [20], and also in forestry [21–23] have been aware of this relationship long before.

In this paper, we use the following terminology that corresponds to the mathematical explanation of derivatives and integrals in calculus [24]: If a state variable, y , is measured at times t_0 and t_1 , or size at both times, we have a dynamic variable, therefore we also refer to them as growth at times t_0 and t_1 (Fig. 1).

The difference between these two measurements is the *increment* of that variable, for the period of time $\Delta t = t_1 - t_0$, and is symbolized by $\Delta y = y_1 - y_0$. The ratio $\Delta y / \Delta t$, is referred to as the *average rate of change*, between t_0 and t_1 , which is how fast the y variable is changing, and is the slope of a straight line connecting $f(t_0)$ to $f(t_1)$ in Fig. 1. If Δt becomes infinitesimally small, then we are analyzing the derivative of y , dy/dt , in y for any particular time (i.e., the slope of the tangent line at points t_0 and t_1 in Fig. 1), which we call (*instantaneous*) *growth rate*. If the increment

in the state variable from t_0 up to a time t_1 , Δy (or dy/dt integrated), is added to y_0 , then we get y_1 , the growth at time t_1 .

If we were able to capture the entire lifetime of most live organisms, their empirical growth would depict a sigmoid curve. A sigmoid or S-type curve resembles trends in the life cycle of many living organisms and phenomena [25] (Fig. 2a). Virtually all exterior tree dimensions develop in a sigmoid manner with respect to time [26]. A growth curve is typically separated in phases (Fig. 2a). Generally three phases have been recognized [27–29], but the pattern remains the same. Baker [28] called them acceleration, intermediate, and deceleration and Assmann [29] youth, full vigor, and old age; while Zedaker et al. [30], youth, maturity, and senescence. In phase 1, y is an exponential function of time [27]. Trees grow relatively slowly at first, increasing their growth-rate to the point of inflection of the growth curve [30]. In phase 2, y is directly proportional to time [27], and in phase 3, there is a decrease of the growth-rates [27, 30].

In addition to *CAI*, the forestry literature recognizes the term *mean annual increment (MAI)*, which is found by dividing the growth through time t by the number of years required to produce it (red curve in Fig. 2b), i.e., $MAI = y_t / t$. Growth and increment curves have long been used for making silvicultural decisions, using as reference the following facts. The inflection point of the *CAI* curve, which occurs at the time symbolized by t^a in Fig. 2, is also the time of maximum *CAI*. It has been recognized as a good opportunity to apply thinning, since the increment of





Fig. 1 Growth definitions for the state variable y obtained from function f . Growth curve (blue), state variable at time 0 ($f(t_0)$) and time 1 ($f(t_1)$), increment (Δy) between t_0 and t_1 , and growth-rate or derivative (dy/dt) at t_0 ($f'(t_0)$) and t_1 ($f'(t_1)$). The derivative, or instantaneous growth-rate, is the slope of the tangent lines (black) at points t_0 and t_1

Fig. 2 Growth and growth-rates curves. **a** The trajectory of the state variable (y), generally representing an expression of size, is the growth curve. Meanwhile, **b** growth-rates curves are represented by the current (blue) and mean (red) annual increments. t^a is the time of culmination of current annual increment and t^b is the time of maximum mean annual increment



residual trees is going to reach its maximum. The point t^a is called the culmination of current annual increment [29–31]. The inflection point of the MAI curve, which occurs at the time symbolized by t^b in Fig. 2, represents the point where the curves of MAI and CAI cross, and MAI is maximized. It has been recognized as a biological criterion to choose the rotation age of even-aged stands. More specifically, it is the rotation that maximizes the growth. It would be the rotation that leads to maximum carbon sequestration if the carbon of harvested biomass would never be released back to the atmosphere [32]. Otherwise, the optimal rotation for carbon sequestration would be longer. Point t^b would be an economically optimal rotation if the interest rate of the forest owner is 0%. A higher interest rate would make the optimal rotation shorter [33].

Growth Equations

Many empirical and theoretical growth equations have been used in forest research. Traditionally, practitioners want to predict the value of a random variable at time t , let's say y_t , as a function of some variables that might affect the value of y_t . The usual approach is to build a mathematical model that fits the pattern of the observed data. The resulting model is called “empirical”. On the other hand, growth

equations have been developed from biological theory about the growth process, with parameters that have (sometimes) biological meaning. Theoretical growth equations have become preferred in research but not always in practice. Some of the so called theoretical growth equations have empirical bases. Nevertheless, we prefer to call them as such because the parameters have some biological interpretation.

Empirical models serve a different purpose than do theoretical models. Thereupon, empirical models should not be viewed as poorer alternatives.

Theoretical growth equations have been developed for many biological disciplines. Several mathematical equations are based on theoretical constructs. Since many growth curves are non-linear in terms of their parameters, growth models are an important family of non-linear models [34]. There is an extensive number of growth equations in the literature; however, it is hard to believe that a model with three or four parameters could describe so complicated process as growth from birth to death [35, 36]. The origin of many non-linear models in use today can be traced to scholarly efforts to discover laws of nature, to reveal scales of being, and to understand the forces of life [34]. There are several growth equations with theoretical or theoretical-empirical bases. Here, we focus on some that have been widely used in forestry research.

There are many different parameterizations of a certain growth model. Because these models start from differential equations, many trajectories can be obtained; as a result numerous models have been proposed based on the differential forms. Furthermore from different differential equations it is possible to attain the same solution or trajectory, a mathematical fact that is not fully understood, as expressed by [37, 38]. Several authors have reported different parameterization of growth models for forestry and other scientific disciplines [3, 34, 37, 39–44]. Here, we present differential forms and solutions of selected and most widely use models in forest science. We want to stress that by selecting these mathematical functions, we are aiming to compare their mathematical differences and origins, but not providing a detailed review of all the available growth functions that can be found.

Hereafter, all the parameters are positive, as well as time (t) is positive, and the state variable y . We also provide expressions of the asymptote (i.e., the maximum level for the state variable) and the points of inflection (i.e., where the curve changes of curvature) of all models. We shall use Greek letters to refer to parameters and italics font style to refer to variables. The use of the same symbols that represent parameters in equations through the document do not represent the mathematical equivalence, unless that is clearly specified. For example, β or γ can be used in different equations, but they do not imply the same number.

Monomolecular Growth Model The monomolecular model is the following differential equation

$$\frac{dy}{dt} = \beta(\alpha - y), \tag{1}$$

where y is the state variable, t represent time, and β and α are parameters that represents the proportional constant and final size, respectively. For this model growth is proportional to the remaining size of the organism [34]. Therefore, the growth-rate decreases as t increases and the equation cannot describe sigmoidal growth. A solution of (1) produces the growth function

$$y_t = \alpha \left\{ 1 - \left(1 - \frac{y_0}{\alpha} \right) e^{-\beta(t-t_0)} \right\}. \tag{2}$$

If we further assume that $t_0 = 0, y_0 = 0$, then (2) becomes

$$y_t = \alpha \{ 1 - e^{-\beta t} \}, \tag{3}$$

where α is the asymptote of the state variable, but the model does not have an inflection point (Table 1). The equation form of (3) is known as the Mitscherlich law or Mitscherlich equation [34].

This function was proposed for height-diameter modelling by [45] as well and is therefore sometimes called Meyer’s equation in the literature.

Table 1 Features of the studied growth models

Model	Differential form	A solution	Asymptote	Point of inflection
Monomolecular	$\frac{dy}{dt} = \beta(\alpha - y)$	$y_t = \alpha(1 - e^{-\beta t})$	α	None
Logistic	$\frac{dy}{dt} = \beta y(1 - \frac{y}{\alpha})$	$y_t = \frac{\alpha}{1 + e^{\beta_0 - \beta t}}$	α	$[\frac{\beta_0}{\beta}; \frac{\alpha}{2}]$
Gompertz	$\frac{dy}{dt} = \beta y(\ln \alpha - \ln y)$	$y_t = \alpha e^{-e^{-\beta(t-\gamma)}}$	α	$[\gamma; \frac{\alpha}{e}]$
Johnson-Schumacher	$\frac{dy}{dt} = \beta d \left(\frac{1}{t} \right)$	$\ln y_t = \alpha - \beta \left(\frac{1}{t} \right)$	e^α	$[\frac{\beta}{2}; e^{\alpha-2}]$
Bertalanffy-Richards	$\frac{dy}{dt} = \left(\frac{\beta}{\gamma} \right) y \left[\left(\frac{\alpha}{y} \right)^\gamma - 1 \right]$	$y_t = \alpha \left[1 - e^{-\beta t} \right]^{\frac{1}{\gamma}}$	α	$[-\frac{\ln(\gamma)}{\beta}; \alpha(1 - \gamma)^{\frac{1}{\gamma}}]$
Weibull	$\frac{dy}{dt} = \alpha \beta \gamma t^{\gamma-1} e^{-\beta t}$	$y_t = \alpha(1 - e^{-\beta t^\gamma})$	α	$\left[\left(\frac{\gamma-1}{\gamma\beta} \right)^{\frac{1}{\gamma}}; \alpha(1 - e^{-\gamma})^{\frac{1}{\gamma}} \right]$
Schnute	$\frac{d^2y}{dt^2} = \frac{dy}{dt} [-\alpha + (1 - \beta)\gamma]$	$y_t = \left\{ y_1^\beta + (y_2^\beta - y_1^\beta) \frac{1 - e^{-\alpha(t-t_1)}}{1 - e^{-\alpha(t_2-t_1)}} \right\}^{\frac{1}{\beta}}$	$\alpha \left[y_1^\beta + \frac{y_2^\beta - y_1^\beta}{1 - e^{-\alpha(t-t_1)}} \right]^{\frac{1}{\beta}}$	$t_1 + t_2 - \frac{1}{\alpha} \ln \left(\frac{\beta(y_2^\beta e^{\alpha t_2} - y_1^\beta e^{\alpha t_1})}{y_2^\beta - y_1^\beta} \right); (1 - \beta) \left(\frac{y_2^\beta e^{\alpha t_2} - y_1^\beta e^{\alpha t_1}}{e^{\alpha t_2} - e^{\alpha t_1}} \right)$

The differential form (i.e., growth-rate equation), a solution (i.e., growth equation), the asymptote and the inflection point are provided

Logistic Growth Model The logistic growth is mostly attributed to Verhulst’s work in 1838 [46–48]. The logistic model imposes a restriction to the exponential growth, then representing limiting resources for population growth.

There are many parametrizations of the logistic model, but we shall show the one most commonly cited in the ecological literature [46, 49],

$$\frac{dy}{dt} = \theta y - \phi y^2 = y(\theta - \phi y), \tag{4}$$

where y is the state variable, θ and ϕ are parameters.

The differential logistic equation can also be parameterized as follows [48, 50], based on treating θ as β and (θ/ϕ) as α ,

$$\frac{dy}{dt} = \beta y - \left(\frac{\beta}{\alpha}\right) y^2 = \beta y \left(1 - \frac{y}{\alpha}\right), \tag{5}$$

where the constant α has a biological interpretation as the carrying capacity of the environment, e.g., the maximum population size for ecological studies or the maximum tree size, and β is a proportionality constant.

The logistic growth equation is the simplest equation describing sigmoidal population growth in a resource-limited environment, and it forms the basis for many models in ecology [48]. A solution of the differential form of the logistic is the following growth curve,

$$y_t = \frac{\alpha}{1 + [(\alpha - y_0)/y_0]e^{-\beta t}}, \tag{6}$$

where y_0 is the value of the initial condition of the state variable y . From (5) we see that when $y \rightarrow \alpha$, which happen when $t \rightarrow \infty$, the growth-rate is zero, therefore the population being modelled or the size of any organism, does not grow without limit. The logistic equation is symmetric around the inflection point t_0 , implying that the growth rate fulfills $f(t_0 - t) = f(t_0 + t)$ for any positive t . Also notice that, when the growth-rate of the equation (Eq. 4) is plotted as a function of the state variable y , and not time, the point of inflection $y = \alpha/2$ (Table 1) is when the growth-rate reaches its maximum.

A usual parameterization found in the forestry literature [31, 51] is

$$y_t = \frac{\alpha}{1 + e^{\beta_0 - \beta_1 t}} \tag{7}$$

which is equivalent to make $(\alpha - y_0)/y_0$ of Eq. (6) as equal to e^{β_0} , and β_0 and β_1 are parameters. Robertson [52] proposed the same differential equation (as in Eq. 4) to describe an autocatalytic monomolecular reaction in biochemistry. Then, the logistic equation is also known as the expression that represents the “autocatalytic law” of physiology and chemistry. When [53] commented on some recent studies on growth including the one by [52], further generalized this thought to brain (and other) growth. According to [34], Pearl and Reed promoted the autocatalytic concept not only for individual but also for

population growth, in their work of 1924. In sociology, the logistic model is know as the law of Verhulst established in 1838 describing the growth of human populations with limited resources [54].

Gompertz Growth Model This growth model was proposed by [55], who was concerned with modelling mortality in an arithmetical progression; however, [56] derived it as a growth model. Gompertz’s differential model is

$$\begin{aligned} \frac{d \ln y}{dt} &= \beta(\ln \alpha - \ln y), \\ \frac{1}{y} \frac{dy}{dt} &= \beta(\ln \alpha - \ln y), \\ \frac{dy}{dt} &= \beta y(\ln \alpha - \ln y), \end{aligned} \tag{8}$$

when the relative growth-rate declines with $\ln y$ and the other terms are parameters.

The most common parameterization of a solution of (8) is [3, 34]

$$y_t = \alpha e^{-e^{-\beta(t-\gamma)}}, \tag{9}$$

where α is the asymptote and γ is the time where the point of inflection occurs (Table 1). Although β is a dimensionless parameter, it affects the maximum growth-rate (i.e., $\alpha\beta/e$). As opposed to the logistic model, the Gompertz curve is asymmetric.

Johnson-Schumacher Growth Model This model was independently proposed by [57] and [21], which has been also known as the “reciprocal function”.

Schumacher [21] explained the model by saying that the growth percent varies inversely with age, which he expressed in a differential form as follows:

$$\frac{dy}{y} = \beta d\left(\frac{1}{t}\right). \tag{10}$$

A solution to this differential equation is found by first, set $u = t^{-1}$ and then substitute $du = -t^{-2}dt$ for $d(1/t) = -t^{-2}dt$ in Eq. (10)

$$\begin{aligned} \frac{dy}{y} &= \beta du, \\ \int \frac{1}{y} dy &= \int \beta du, \\ \ln y + \gamma_1 &= \beta u + \gamma_2, \end{aligned} \tag{11}$$

where γ_1 and γ_2 are the constants of integration. Second, by assigning $\alpha = \gamma_2 - \gamma_1$, we arrive at

$$\ln y = \alpha + \beta \left(\frac{1}{t}\right). \tag{12}$$

This model has been widely used in forestry, mainly for fitting height-age models. Equation (12) allows parameter estimation in the context of linear modelling. However, one should notice that model fitting for $\ln y$ leads to biased

predictions of y [58, 59]. Also the alternative form $y = \alpha \exp(\frac{\beta}{t})$, is used in forest sciences; however, here α has a different interpretation than in (12).

The differential equation of [21] can be represented as the growth-rate as a fraction of the state variable y , as follows

$$\left(\frac{dy}{d\left(\frac{1}{t}\right)}\right)/y = \beta. \tag{13}$$

however, as mentioned by [60], this differential equation does not represent what [21] said in words, i.e., “growth per cent varies inversely with age”. Instead, that thought is represented by the following differential equation

$$\left(\frac{dy}{dt}\right)/y = \beta \left(\frac{1}{t}\right). \tag{14}$$

A solution of Eq. (14) is obtained as follows,

$$\begin{aligned} \frac{dy}{y} &= \beta \left(\frac{1}{t}\right) dt, \\ \int \frac{1}{y} dy &= \beta \int \left(\frac{1}{t}\right) dt, \\ \ln y + \gamma_1 &= \beta \ln t + \gamma_2, \end{aligned} \tag{15}$$

where γ_1 and γ_2 are the constants of integration, and $\alpha = \gamma_2 - \gamma_1$, as before, we get

$$\ln y = \alpha + \beta \ln t, \tag{16}$$

This function is not the growth equation proposed by [21], which is being advocated to represent growth-rate as a percentage inversely proportional to time. Using exponential transformation, we see that the equation is a linearized version of the extensively used power function $y = \phi t^\alpha$, where $\phi = e^\alpha$.

Furnival [60] also detected the error in [21], and said “the curve given by [21] is obtained if growth percentage is taken as inversely proportional to the square of age”. If we proceed as above, we arrive to

$$\ln y = \alpha + \beta \left(-\frac{1}{t}\right). \tag{17}$$

This formulation emphasizes that model (12) leads to increasing logarithmic growth only if $\beta < 0$, whereas in (17) we need to have $\beta > 0$. Therefore, we favor this expression as a correct solution of the differential equation

$$\frac{dy}{y} = \beta d\left(\frac{1}{t^2}\right). \tag{18}$$

Finally, we could say that the derivation of Schumacher is correct but his formulation is wrong.

Bertalanffy Growth Model This model was first published in German [61], and later the author introduced it to the English literature in a brief paper [62], and in a more developed one [20].

He originally developed it for weight growth of animals. The model is a result of a deeper developed basic idea of Pütter in 1920 [54]. That is, growth can be considered a result of a counteraction of synthesis and destruction, of the anabolism and catabolism of the building materials of the body. The differential form of the von Bertalanffy model is

$$\frac{dy}{dt} = \eta y^\theta - \kappa y^\phi, \tag{19}$$

where η and κ are constants of anabolism and catabolism respectively, and the exponents θ and ϕ indicate that both are proportional to some power of the state variable y . Leary [63] termed (19) as the Bertalanffy’s anabolic-catabolic balance equation of biological growth. As explained by [20], this differential equation represents “the change of y is given by the difference between the processes of building up and breaking down”. Nonetheless, this is a property that is also shared by the monomolecular (1), logistic (4) and Gompertz (8) equations, in the sense that the differential equation can be separated into different additive terms. In the Bertalanffy model, both these processes are described by the simple power models, but the form of these submodels is not justified by biological theory. Ricker [47] pointed out that von Bertalanffy did not make clear what he meant by anabolism and catabolism, and aside from all the widely quoted references to his model as a solid theoretical growth model, that is not actually true. For instance, von Bertalanffy [54] explained, when referring to his model, that the use of the exponent θ and ϕ in (19) has a pure empirical base, since it is well known that the size dependence of physiological processes can well be approximated by allometric expressions. More recently, Seber and Wild [3] noticed that unfortunately the biological basis of these models (referring to the logistic, Gompertz, and von Bertalanffy) has been taken too seriously and had led to the use of ill-fitting growth curves.

Bertalanffy [20] justified that $\phi = 1$ based on physiological experience; hence

$$\frac{dy}{dt} = \eta y^\theta - \kappa y, \tag{20}$$

and he also empirically found that for a wide class of animals, the allometric power for the metabolic rate (θ) is 2/3. Thus, differential equation (19) becomes

$$\frac{dy}{dt} = \eta y^{2/3} - \kappa y. \tag{21}$$

Equation (21) is usually termed the von Bertalanffy model [3]; nonetheless, we prefer to reserve that denomination to Eq. (20).

Bertalanffy-Richards Growth Model Richards [19] doubted some theoretical considerations of the Bertalanffy model (21), and treated some of the parameters to be related to a

point of inflection [3]. The Richard’s differential equation is

$$\frac{dy}{dt} = \left(\frac{\beta}{\gamma}\right) y \left[\left(\frac{\alpha}{y}\right)^\gamma - 1 \right], \tag{22}$$

where α, β and γ are the asymptote, a scale, and shape-related parameters [34]. Although, this equation has been termed as the Bertalanffy-Richards model by [64, 65], and also as a generalization of the Bertalanffy model [19], it is paramount to understand that it is simply a reparametrization of (20) as shown by [38]. A solution of this differential equation is

$$y_t = \alpha \left\{ 1 - \left[1 - \left(\frac{y_0}{\alpha}\right)^\gamma \right] e^{-\beta(t-t_0)} \right\}^{\frac{1}{\gamma}}. \tag{23}$$

Notice that (23) becomes the following growth equation if $y_0 = t_0 = 0$,

$$y_t = \alpha \left\{ 1 - e^{-\beta t} \right\}^{\frac{1}{\gamma}}. \tag{24}$$

Hence, (24) is a special case of (23), being the former suitable for forestry plantations and the latter for natural forests [66].

It is key to realize, and as aforementioned, that from distinct differential equations, we can arrive at a same solution. Notice that from the following linear differential equation with a power transformation [64],

$$\frac{dy^\gamma}{dt} = \beta (\alpha^\gamma - y^\gamma), \tag{25}$$

we can obtain the same solution of (23), as explained in detail by [38].

Chapman [67] starting from the von Bertalanffy differential equation, in the same way as [19] did, derived a new parameterization of the model particularly suitable for fisheries population modelling. Because of that, sometimes this equation has been called “Chapman-Richards” as well [40, 68]. We can argue that the denomination for the generalized differential equation (20) should be credited to [20], because even though he did not state it formally in his paper, he used it as the basis to obtain his proposed solution. Nevertheless, Richards [19] made this clearer, and also gave his famous 2/3 exponent for θ in (20). Thus, the denomination “Bertalanffy-Richards” seems a better compromise. Any other denomination, e.g., Chapman-Richards, even though correct in the sense that they started from the differential equation (20), is burdensome. Similar claims had been brought by [37–39, 69]. Based on the shortcomings mentioned above, we shall focus on the Bertalanffy-Richards growth model (23).

Weibull Growth Model This model is a probability density function (pdf) widely used in engineering [70] as well as in forestry for describing diameter distributions [71, 72] because of its versatility. Prodan [73] noticed that tree increment and growth curves resemble a probability density

(i.e., frequency distribution) and a probability distribution (i.e., cumulative frequencies distribution), respectively. Later, Yeng et al. [11] exploited this relationship and used it to fit growth curves of trees. Based on the review given by [74], it seems that [11] were the first to notice this. The differential form of the Weibull growth model is as follows [40]:

$$\frac{dy}{dt} = \alpha \beta \gamma t^{\gamma-1} e^{-\beta t^\gamma}. \tag{26}$$

A solution of the Weibull growth-rate model is [11, 41]

$$y_t = \alpha \left\{ 1 - e^{-\beta t^\gamma} \right\}, \tag{27}$$

where α, β , and γ are parameters representing the asymptote, intrinsic rate and the shape of the curve, respectively. The last two parameters affects the point of inflection (Table 1). Zeide [40, 41] and Payandeh and Wang [75] found that the Weibull growth model performed as well as other common growth equations in modelling tree growth of several variables.

Schnute Growth Model Schnute [76] introduced a model first tested for fisheries research. The model is capable of assuming a wide range of characteristic shapes that describe asymptotic as well as non-asymptotic growth trends [65, 77]. The Schnute’s model is based upon acceleration in growth, that is the rate of a rate, hence using second derivatives as follows [76]:

$$\frac{d^2y}{dt^2} = \frac{dy}{dt} \left[-\alpha + (1 - \beta) \frac{1}{y} \right], \tag{28}$$

where α, β , and dy/dt are parameters. A solution of the Schnute differential equation is [76]

$$y_t = \left\{ \tau_1^\beta + (\tau_2^\beta - \tau_1^\beta) \frac{1 - e^{-\alpha(t-t_1)}}{1 - e^{-\alpha(t_2-t_1)}} \right\}^{1/\beta}, \tag{29}$$

where $\alpha \neq 0, \beta \neq 0, t$ is the time of interest, t_1 time at beginning (e.g., young), t_2 time at end of (e.g., old), τ_1 and τ_2 are parameters representing the state variable y at time t_1 and t_2 , respectively. As stated by [65], ϕ is a constant acceleration in growth-rate, and β is incremental acceleration. Unfortunately, the growth function (29) does not have a single parameter with biological interpretation (Table 1). The Schnute growth-rate model is among the very few growth investigations based upon second derivatives, with the exception of the lucid study of [78] when modelling height growth.

The Use of Growth Equations in Empirical Studies

Although the logistic model has been long used in population dynamics in ecology, it has barely been used in forestry to model growth [51]. Among them, Cooper [22],

MacKinney and Schumacher [31], and Monserud [51] used it for modelling growth of stand volume, stem area, and tree height growth, respectively.

Unlike the logistic model, the Gompertz model is not symmetric about the point of inflection.

It is a model more used in ecology than in forestry; it performed well in population models as reported by the comparison of [79] and was used with success by [80]. In forestry, it was first mentioned by [81], but was not used until the work of [82] in modelling stand volume growth.

The Bertalanffy-Richards model is used more than any other function in studies at tree and stand level [41, 43].

This equation was first used in forestry by [83], but mostly popularized by [68], after which the model was termed and later known as ‘‘Chapman-Richards’’.

The Schumacher growth equation (17) is long-established for modelling height growth of dominant trees, because it is linear in its parameters. Notwithstanding, lately the Bertalanffy-Richards model has become the preferred model.

The Weibull model was first proposed by [11]. Prodan [73] earlier had mentioned the resemblance of growth and increment curves to probability density and probability distribution.

The Schnute model has the property of having asymptotic as well as non-asymptotic shape depending on the parameter values combination, therefore being versatile to a wider ranges of situations. Bredenkamp and Gregoire [77] were

the first to introduce the model to forestry. Later on, Yuncai et al. [65, 84] and [85] used this model as well.

Comparisons among growth equations are valuable both for practical and theoretical use; nonetheless, it seems that researchers tend to prefer one of the models beforehand.

Researchers might conduct initial screenings, but may not report the results in detail. Considering the large number of growth functions [43], very few studies have empirically compared them. In Table 2, we have summarized a comprehensive list of studies in this regard. Overall, the authors selected their models using traditional goodness-of-fit statistics. The Bertalanffy-Richards, Weibull and Schnute were the most widely selected according to these criteria.

Practical Demonstration

To illustrate the different growth-rates, we estimated the parameters of the growth equations by using time series height data of 107 *Nothofagus dombeyi* sample trees, a native evergreen tree species of South America temperate forests [93], growing on secondary stands in the south of the country. The tree-level data were collected within 30 sample plots established throughout the ecological distribution of the species. Sample trees, at the time of sampling, ranged from young ages to mature (between 21 and 71 years), as well as from small trees to taller ones (between 10 and 34 m, Table 3).

Table 2 Empirical comparisons of growth equations in some forestry studies

Source	Model							Data
	Monomolecular	Logistic	Gompertz	Bertalanffy-Richards	Weibull	Schnute	Other	
[11]		•	•	•	✓			Tree volume
[86]	✓		•				•	Stand volume
[77]				•		✓		Tree diameter
[40]		•	•	•	•		✓	Tree diameter
[75]				•	✓			Tree height
[87]	•	•	•		✓			Plant disease
[65]				✓		✓		Tree diameter
[88]			•	✓	✓	✓		Tree height
[84]				✓		✓		Tree height
[89]		•	•	✓	✓	✓		Tree height
[43]	•	•	•	•	✓			Tree height
[90]				✓	•			Tree basal area
[91]	•	✓	•	•		•	•	Weight
[85]						✓	•	Tree height
[92]				✓	•		✓	Plant disease

The symbol • and ✓ represents that the equation was assessed and selected as the best in the study, respectively

Table 3 Tree level variables summary for the 107 *Nothofagus dombeyi* trees at the time of sampling. *d* is diameter at breast height, *h* is total height, *age* is total age, and *bha* is breast-height age

Statistic	<i>d</i> (cm)	<i>h</i> (m)	<i>age</i> (yr)	<i>bha</i> (yr)
Minimum	5.3	9.9	21	15
Maximum	60.2	33.7	71	68
Mean	26.6	19.9	41.3	37.7
Median	26	20.8	40	37
CV(%)	19	16.6	15.8	17.3

The height growth data cover a wide variety of growing conditions (Fig. 3). Further details on the data can be found in [38].

The growth equations were fitted in a mixed-effects model framework for considering the grouped structure of the data [66, 94].

A general non-linear mixed-effects model for the *k*th observation on the *j*th tree at the *i*th plot is

$$y_{ijk} = f(\phi_{ij}, t_{ijk}) + \varepsilon_{ijk}, \tag{30}$$

where *f* is a growth equation and the parameter vector ϕ_{ij} includes the parameters of the selected growth function. Each parameter consists of a fixed effect (μ) and two random effects, i.e., plot (b_i) and tree (b_{ij}), therefore

$$\phi_{ij} = \mu + b_i + b_{ij}. \tag{31}$$




Fig. 3 Tree height growth series of dominant trees for *Nothofagus dombeyi* in southern Chile. Gray lines join successive observations of height on the same tree

We assume that the random effects among plots are independent with $b_j \sim N(\mathbf{0}, \mathbf{D})$, as well as for the random effects among trees within plots with $b_{ij} \sim N(\mathbf{0}, \mathbf{E})$. Also the residual errors are independent and normally distributed with $\text{Var}(\varepsilon_{ijk}) = \sigma^2$. Further details on the formulation of non-linear mixed-effects models are presented in [59, 95–97].

The models were fitted using the nlme package [98] of R [99].

The fixed-effects parameter estimates for each growth model are displayed in Table 4. We also computed the prediction statistics as in [100], i.e., the root mean square differences (RMSD) and the aggregated difference (AD) using the population level predicted value from the corresponding mixed-effects fitted model. These statistics provide some insight into model fit. Except for the Johnson-Schmacher equation, all models have similar precision and accuracy levels.

Using the fixed-effects estimated parameters of the non-linear mixed-effects growth models (Table 4), we represent the height growth or trajectory of the state variable *h* (Fig. 4a), and the instantaneous growth-rates versus time (Fig. 4b) and versus the state variable (Fig. 4c), as well. These trajectories should be interpreted as the estimates for a typical sample plot of the data set.

The number of parameters (i.e., coefficients) affects the flexibility of functions. As such, the greater number of parameters, the greater flexibility or intrinsic curvature. We used two functions (i.e., Monomolecular and Johnson-Schumacher) having 2 parameters, four (i.e., logistic, Gomperts, Bertlanffy-Richards, and Weibull) having 3 parameters, and the Schnute’s function having 4 parameters.

Comparing flexibility of the functions for a given number of parameters is of great interest when a generally applicable function is searched for empirical growth modelling. See [101] for such an analysis in the context of diameter distribution models. However, we leave such analysis for future studies.

Caveats

In physics, it is rare to analyze different models because they are already well established, and uncertainty is not a central feature. On the contrary, in biology-related disciplines, such as forestry, variability and uncertainty is the rule. As such, selecting a growth equation on which our findings are based is not straightforward. Most practitioners might compare a suite of equations and choose one of the merit of goodness-of-fit indices [102, 103]. On the contrary, others will start with a preferred growth equation at the very beginning of their studies [104–106].

In this vein, it is often not recognized the tremendous impact that a selected growth model will have on our

Table 4 Fixed-effects parameter estimates and prediction statistics for each mixed-effects growth model. All models were fitted by maximum-likelihood and using time series height data of 107

Nothofagus dombeyi sample trees from southern Chile. RMSD is the root mean square differences and AD the aggregated difference

Growth model	Parameters			RMSD		AD		
	$\hat{\alpha}$	$\hat{\beta}$	$\hat{\gamma}$	(m)	%	(m)	%	
Monomolecular	49.76	0.012	–	2.8249	26.2	0.0507	0.47	
Logistic	23.41	2.242	0.10	2.8560	26.5	0.0689	0.63	
Gompertz	26.28	0.057	18.59	2.8003	25.9	0.1082	1.00	
Johnson-Schumacher	2.72	5.681	–	5.1607	47.9	1.6846	15.62	
Bertalanffy-Richards	34.67	0.024	0.81	2.7876	25.9	0.1494	1.39	
Weibull	32.70	0.012	1.18	2.7871	25.8	0.1487	1.38	
Schnute	0.16	-2.13	$\hat{\tau}_1$ 2.84	$\hat{\tau}_2$ 18.02	2.9419	27.3	0.2120	1.96

inferences. For instance, let us assume that the goodness-of-fit indices are similar among the suite of growth equations examined here as a reference, a pattern that is not unrealistic. However, if we are interested in finding the maximum height for the species, a critical functional trait, the asymptote of the growth curves (Fig. 4a) varies tremendously between an unrealistic value of 15.2 m for the Johnson-Schumacher model (we used the [58] bias correction for prediction) and 49.8 m for the monomolecular one (Table 4).

Overall, there is no theoretical reason to limit either tree or forest growth analysis by using a single equation, and claims on different reparametrizations have been proposed [107].

As previously, most growth models are non-linear. The only linear model (in the parameters) assessed here was the Johnson-Schumacher (Eq. 17). There is no closed formula to estimate non-linear models' parameters, as we have for the linear models. Nowadays, most statistical software

includes tools for easing the fit of non-linear equations, such as self-starting functions that find good initial guesses for the parameters [97]. However, a little more care and time may be needed than with linear models, e.g., to run the numerical estimation routines in large data sets, to find good enough initial guesses for the parameters, and to ensure that the estimates found are the global maximum of the likelihood function, but not local [3].

Regardless, several attempts have been devoted to ecological applications to offer guidance for fitting non-linear growth models, since [108, 109] to [44], highlighting that this is still an important topic when modelling tree and forest growth. Consequently, the possibility of working with models having parameters with biological or physical interpretation is an advantage for fitting these non-linear models.

There are numerous applications for the unified mathematical treatment of growth equations presented here. The




Fig. 4 Growth (a) and growth-rates (b) versus time, and growth-rates versus the state variable c for the growth equations studied here

more obvious are in forest modelling, such as the ones fully described in [110, 111]. Besides, growth equations have several implications in decision support and scientific findings [112]. Nonetheless, there are a handful of other applications where growth equations are used. Among them, we can mention the studies focus on: ecophysiological aspects of sapling growth [104–106, 113]; forest-dynamics [114]; silviculture [115, 116]; and restoration ecology [117–119], where all the findings are based upon a single fitted-and-assessed growth function.

As lucidly summarized by [3], we can use x (i.e., another state variable) instead of t in the growth-rate equations, and by this simple twirl, the same mathematical models shown here, as well as their implications, can serve for representing tree allometry [120], a major scientific endeavor in biology [121, 122]. In the forestry context, growth equations have been used in modelling height-diameter [88, 123, 124], and stem taper [125, 126] allometric relationships.

Growth-rates of the tree and forest growing processes are essential in research [23, 127, 128], especially to study the partitioning of components, resources, and allocation of them to different processes. At least a basic but correct mathematical understanding of differential equations is paramount for researching these complex processes. Similar advocacy has been claimed by others in ecology [129]. It is noteworthy to realize how the forestry literature is way more focused on growth than on the growth-rates themselves; that is to say, forestry is a discipline where we seem to be trained to think in an integrated form rather than in a differential form. In contrast, we highlight the pioneer works on differential equations in forestry led by [64, 130–133].

Although we might think that the reasons for focusing on growth, but not on growth-rates when modelling forestry variables are a handful; the main one could be related to the fact that instantaneous growth rates are essentially unobservable, whereas realized sizes can be measured and so are amenable to exploratory data analysis. However, this can also be applied to other biological disciplines, such as wildlife ecology and landscape ecology. Regardless, the use of growth-rates, or differential equations, are just a few compared to size-time equations (i.e., a differential equation solution). Maybe the reasons are also related to a weak educational background in calculus of ecology-related scientists [129, 134]. Accordantly, we favor using growth-rates functions, such as differential equations, in forest growth modelling when justified by the research question [23, 38, 128, 133].

As a reflection of the above, the analyses of the growth rates is mandatory. The growth-rates against time for each model broadly vary (Fig. 4b). All of them implies completely different processes. For instance, the Monomolecular and the Johnson-Schumacher curves seem

unrealistic. Both the Bertalanffy-Richards and the Weibull functions behave quite similar at reaching their maximum growth-rates at much earlier stages. On the contrary, the logistic, Gompertz and Bertalanffy-Richards seems to be more feasible. Finally, a graphic representing the growth-rates against time (Fig. 4b) is usually the most commonly reported; however, plotting the growth-rates against size (Fig. 4c) is also essential.

These two figures describe completely different processes. For instance, the logistic growth-rate is symmetric, and the point on which this occurs is where half of the asymptote is reached, and the growth-rate is maximum. Nonetheless, no biological reasons for the symmetry have been presented. The asymmetric behavior of the other functions seems more realistic. Both the Bertalanffy-Richards and the Weibull behave quite similarly, reaching their maximum growth-rates at much smaller sizes. Finally, the Monomolecular and the Johnson-Schumacher curves are entirely unrealistic.

In general, the rules on which the growth equations are based are rather broad and motivated more by mathematical formulation than by a solid, established theory about the biological processes of growth.

Concluding Remarks

We have shown that the mathematical differences among several growth equations can be more easily explained and understood using their differential equation than their integrated forms. Inter alia, the assumed-and-claimed biological basis of these growth-rate models has been taken too seriously, and the focus should be on using a plausible equation for the organism being modelled. More attention should be put on parameter estimation strategies and behavior analysis of the proposed models. It is difficult for a single function to capture all possible shapes and rates that such a complex biological process as tree growth can depict in nature. Therefore, comparisons of some functional forms (at least simple ones) must be carried out before selecting a function for drawing scientific findings.

Funding This study was supported by the Chilean research grants Fondecyt No. 1191816 and FONDEF No. ID19|10421, Academy of Finland Flagship Programme (UNITE, decision number 337655), and ANID BASAL FB210015.

Declarations

Conflict of Interest The authors declare that they have no conflict of interest

Human and Animal Rights and Informed Consent This article does not contain any studies with human or animal subjects performed by the authors.

Papers of particular interest, published recently, have been highlighted as:

- Of importance
- Of major importance

References

1. Seger SL, Harlow SD. Mathematical models from laws of growth to tools for biological analysis: fifty years of growth. *Growth*. 1987;51:1–21.
2. Lambers H, Oliveira RS. *Plant physiological ecology*, 3rd ed. New York: Springer; 2019.
3. Seber GAF, Wild CJ. *Nonlinear regression*. New York: Wiley; 1989, p. 800.
4. Evans GC. *The quantitative analysis of plant growth*. Berkeley: University of California Press; 1972, p. 734.
5. Oliver CD, Larson BC. *Forest stand dynamics*, 2nd ed. New York: Wiley; 1996, p. 520.
6. Barnes BV, Zak DR, Denton SR, Spurr SH. *Forest ecology*, 4th ed. New York: Wiley; 1997, p. 800.
7. Kimmins JP. *Forest ecology*, 3rd ed. Upper Saddle River: Pearson Prentice Hall; 2004, p. 611.
8. Odin H. *Studies of the increment rhythm of scots pine and Norway spruce plants*. Technical report, Studia Forestalia Suecica No. 97, Stockholm, Sweden, pp 32. 1972.
9. Ek AR, Monserud RA. *Methodology for modeling forest stand dynamics*. Department of Forestry, University of Wisconsin–Madison, Staff Paper Series No. 2. Madison, WI, USA, pp 30. 1975.
10. Bruce D. Consistent height-growth and growth-rate estimates for re-measured plots. *Forest Sci*. 1981;27(4):711–725.
11. Yang RC, Kozak A, Smith JH. The potential of Weibull-type functions as flexible growth curves. *Can J For Res*. 1978;8(2):424–431.
12. Buckman RE. *Growth and yield of red pine in Minnesota*. USDA For. Serv. Tech. Bull. No. 1272, USA, pp 50. 1962.
13. Haig IT. *Second-growth yield, stand, and volume tables for the western white pine type*. USDA For. Serv. Tech. Bull. No. 323, USA, pp 67. 1932.
14. • Clutter JL. Compatible growth and yield models for loblolly pine. *Forest Sci* 1963;9(3):354–371. **The author highlights the mathematical relationship between growth and growth-rates in forest growth models.**
15. Curtis RO. A method of estimation of gross yield of Douglas-fir. *For. Sci. Monogr.*, pp 13. 1967.
16. Vanclay JK. *Modelling forest growth and yield: applications to mixed tropical forests*. CAB International, Wallingford, Berkshire, England, pp 312. 1994.
17. Blackman VH. The compound interest law and plant growth. *Ann Bot*. 1919;33(3):353–360.
18. Fisher RA. Some remarks on the methods formulated in a recent article on “the quantitative analysis of plant growth”. *Ann Appl Biol*. 1921;7:367–372.
19. Richards FJ. A flexible growth function for empirical use. *J Experimental Botany*. 1959;10(29):290–300.
20. von Bertalanffy L. Quantitative laws in metabolism and growth. *The Quarterly Review of Biology*. 1957;32(3):217–231.
21. Schumacher FX. A new growth curve and its application to timber yield studies. *J Forestry*. 1939;37(10):819–820.
22. Cooper CF. Equations for the description of past growth in even-aged stands of ponderosa pine. *Forest Sci*. 1961;7(1):72–80.
23. Pommerening A, Muszta A. Relative plant growth revisited: towards a mathematical standardisation of separate approaches. *Ecol Model*. 2016;320:383–392.
24. Larson RE, Hoestetler RP. In: Head DC, editor. *Calculus with analytic geometry*, 3rd ed. Lexington: Company; 1990, p. 1083.
25. Davenport CB. Human growth curve. *J General Physiology*. 1926;10(2):205–216.
26. Monserud RA. *Methodology for simulating Wisconsin northern hardwood stand dynamics*. Ph.D. thesis, University of Wisconsin, Madison, WI, USA, pp 156. 1975.
27. Priestley JH, Pearsall WH. *Growth studies. II. An interpretation of some growth-curves*. *Ann Bot*. 1922;36(2):239–250.
28. Baker FS. *Principles of silviculture*. McGraw-Hill, New York, USA, pp 414. 1950.
29. Assmann E. *The principles of forest yield study*. English edition. Pergamon Press, Oxford, England, pp 506. 1970.
30. Zedaker SM, Burkhart HE, Stage AR. 1987. General principles and patterns of conifer growth and yield. In Walstad, JD, Kuch, PJ eds. *Forest vegetation management for conifer production*. Wiley, New York, USA, p. 203–241.
31. MacKinney AL, Schumacher FX. Construction of yield tables for nonnormal loblolly pine stands. *J Agricultural Research*. 1937;54(7):531–545.
32. Pukkala T. Carbon forestry is surprising. *Forest Ecosystems*. 2018;5(1):11.
33. Viitala E. *Timber, science and statecraft: the emergence of modern forest resource economic thought in Germany*. *Eur J For Res*. 2016;135(6):1037–1054.
34. Schabenberger O, Pierce FJ. *Contemporary statistical models for the plant and soil sciences*. CRC Press, Boca Raton, FL, USA, pp 738. 2002.
35. Koops WJ. Multiphasic growth curve analysis. *Growth*. 1986;50:169–177.
36. Landsberg J, Sands P. *Physiological ecology of forest production: principles, processes and models*. Academic Press, San Diego, CA, USA, pp 330. 2011.
37. García O. Unifying sigmoid univariate growth equations. *Forest Biometry, Modelling and Information Sciences*. 2005;1:63–68.
38. Salas-Eljatib C. Height growth–rate at a given height: a mathematical perspective for forest productivity. *Ecol Model*. 2020;109198:431.
39. Grosenbaugh LR. Generalization and reparameterization of some sigmoid and other nonlinear functions. *Biometrics*. 1965;21(3):708–714.
40. Zeide B. Accuracy of equations describing diameter growth. *Can J For Res*. 1989;19(10):1283–1286.
41. • Zeide B. Analysis of growth equations. *Forest Sci*. 1993;39(3):594–616. **A comprehensive study offering mathematical grounds to compare growth equations.**
42. Shvets V, Zeide B. Investigating parameters of growth equations. *Can J For Res*. 1996;26(11):1980–1990.
43. Kiviste A, Álvarez JG, Rojo A, Ruíz AD. *Funciones de crecimiento de aplicación en el ámbito forestal*. Monografías INIA, Forestal No. 4, Madrid, Spain, pp 190. 2002.
44. Paine CET, Marthews TR, Vogt DR, Purves D, Rees M, Hector A, Turnbull LA. How to fit nonlinear plant growth models and calculate growth rates: an update for ecologists. *Methods Ecol Evol*. 2012;3(2):245–256.
45. Meyer HA. A mathematical expression for height curves. *J Forestry*. 1940;38(5):415–420.
46. Pielou E. *An introduction to mathematical ecology*. New York: Wiley–Interscience; 1969, p. 286.
47. Ricker WE. *Growth rates and models*. Fish physiology. Volume VIII bioenergetics and growth. In: Hoar WS, Randall DJ, and Brett JR, editors. New York: Academic Press; 1979. p. 677–743.

48. Gotelli NJ. A primer of ecology, 3rd ed. Sunderland: Sinauer Associates Inc.; 2001, p. 265.
49. Dennis B, Desharnais RA, Cushing JM, Henson SM, Costantino RF. Can noise induce chaos? *Oikos*. 2003;102(2):329–339.
50. Dennis B. Allee effects in stochastic populations. *Oikos*. 2002;96(3):389–401.
51. Monserud RA. Height growth and site index curves for Inland Douglas-fir based on stem analysis data and forest habitat type. *Forest Sci*. 1984;30(4):943–965.
52. Robertson TB. On the normal rate of growth of an individual, and its biochemical significance. *Archiv für Entwicklungsmechanik der Organismen*. 1908;XXV:581–614.
53. Pearl R. Some recent studies on growth. *The American Naturalist*. 1909;43(509):302–316.
54. von Bertalanffy L. General system theory. foundations, Development, Applications. George Braziller, INC., New York, USA, pp 295. 1969.
55. Gompertz B. On the nature of the function expressive of the law of human mortality, and on a new mode of determining the value of life contingencies. *Philos Trans R Soc Lond*. 1825;115:513–583.
56. Medawar PB. The growth, growth energy, and ageing of the chicken's heart. *Proceedings of the Royal Society B*. 1940;129(856):332–355.
57. Johnson NO. A trend line for growth series. *J Am Stat Assoc*. 1935;30(192):717–717.
58. Baskerville GL. Use of logarithmic regression in the estimation of plant biomass. *Can J For Res*. 1972;3:49–53.
59. ● Mehtätalo, L, Lappi, J. Biometry for forestry and environmental data: with examples in R. Chapman & Hall / CRC Press, Boca Raton, FL, US, pp 411. 2020. **An applied statistics book on forestry data, with emphasis on mixed-effect models.**
60. Furnival GM. Growth and yield prediction: some criticisms and suggestions. *Proceedings of the seminar series predicting forest growth and yield: current issues, future prospects*. College of Forest Resources, University of Washington, Institute of Forest Resources Contribution Number 58. Seattle, WA, USA, pp 23–30. In: Chapell HN and Maguire DA, editors; 1987.
61. von Bertalanffy L. Untersuchungen über die Gesetzmäßigkeit des Wachstums. I. Allgemeine Grundlagen der Theorie. Mathematisch-physiologische Gesetzmäßigkeiten des Wachstums bei Wassertieren. *Wilhelm Roux Arch Entwickl Mech Org*. 1934;131:613–652.
62. von Bertalanffy L. Problems of organic growth. *Nature*. 1949;163:156–158.
63. Leary RA. Interaction theory in forest ecology and management. Martinus Nijhoff/Dr. W. Junk Publishers, Dordrecht, USA, pp 219. 1985.
64. ● García O. A stochastic differential equation model for the height growth of forest stands. *Biometrics*. 1983;39(4):1059–1072. **The author offers a parameter estimation algorithm to fit stochastic differential equations in forestry. Besides, the study provides a plethora of mathematical details.**
65. Yuncai L, Pacheco C, Wolfango F. Comparison of Schnute's and Bertalanffy-Richards' growth functions. *For Ecol Manage*. 1997;96:283–288.
66. Salas C, Stage AR, Robinson AP. Modeling effects of overstory density and competing vegetation on tree height growth. *Forest Sci*. 2008;54(1):107–122.
67. Chapman DG. Statistical problems in dynamics of exploited fisheries populations. *Proc. 4th Berkeley symp. on mathematics, statistics and probability*. In: Neyman J, editors. Berkeley: University of California Press; 1961. p. 153–168.
68. Pienaar LV, Turnbull KJ. The Chapman-Richards generalization of von Bertalanffy's growth model for basal area growth and yield in even-aged stands. *Forest Sci*. 1973;19(1):2–22.
69. García O. Visualization of a general family of growth functions and probability distributions – the growth-curve explorer. *Environ Model Softw*. 2008;23(12):1474–1475.
70. Ross S. A first course in probability, 7th ed. Upper Saddle River: Pearson Prentice Hall; 2006, p. 565.
71. Clutter JL, Fortson JC, Pienaar LV, Brister GH, Bailey RL. Timber management: a quantitative approach. Wiley, New York, USA, pp 333. 1983.
72. Husch B, Beers TW, Kershaw JA. Forest mensuration, 4th ed. New York: Wiley; 2003, p. 443.
73. Prodan M. Forest biometrics. Pergamon Press, London, UK, pp 447. 1968.
74. Pielou EC. The usefulness of ecological models: a stock-taking. *The Quarterly Review of Biology*. 1981;56(1):17–31.
75. Payandeh B, Wang Y. Comparison of the modified Weibull and Richards growth function for developing site index equations. *New Forest*. 1995;9(2):147–155.
76. Schnute I. A versatile growth model with statistically stable parameters. *Can J Fish Aquat Sci*. 1981;38(9):1128–1140.
77. Breckenkamp BV, Gregoire TG. A forestry application of Schnute's generalized growth function. *Forest Sci*. 1988;34(3):790–797.
78. ● Leary RA, Nimerfro K, Holdaway M, Brand G, Burk T, Kolka R, Wolf A. Height growth modeling using second order differential equations and the importance of initial height growth. *For Ecol Manage*. 1997;97(2):165–172. **The study accentuates the effect of initial tree height in the growth trajectories built from differential equations.**
79. Rotella JJ, Ratti JT, Reese KP, Taper ML, Dennis B. Long-term population analysis of gray partridge in eastern Washington. *Journal of Wildlife Ecology*. 1996;60(4):817–825.
80. Dennis B, Ponciano JM, Lele SR, Taper ML, Staples D. Estimating density dependence, process noise and observation error. *Ecol Monogr*. 2006;76(3):323–341.
81. Rawat AS, Franz F. Growth models for tree and stand simulation, Res. Note 30. *Proc. IUFRO Working Party S4.01-4 Meetings, Dep. For. Yield Res., Royal Coll. For., Stockholm, Sweden*, pp 180–221. In: Fries J, editors; 1974.
82. Nokoe S. Demonstrating the flexibility of the Gompertz function as yield model using mature species data. *Commonw For Rev*. 1978;57(1):35–42.
83. Moser JW Jr, Hall OF. Deriving growth and yield functions for uneven-aged forest stands. *Forest Sci*. 1969;15(2):183–188.
84. Yuncai L, Pacheco C, Wolfango F. Features and applications of Bertalanffy-Richards' and Schnute's growth equations. *Nat Resour Model*. 2001;14(3):1–19.
85. Lei Y, Zhang SY. Comparison and selection of growth models using the Schnute model. *J Forest Science*. 2006;52(4):188–196.
86. Leech JW, Ferguson IS. Comparison of yield models for unthinned stands of radiata pine. *Aust For Res*. 1981;11:231–245.
87. Xiao CL, Subbarao KV, Zeng SM. Incorporating an asymptotic parameter into the Weibull model to describe plant disease progress. *J Phytopathology*. 1996;144(7–8):375–382.
88. Zhang L. Cross-validation of non-linear growth functions for modelling tree height-diameter relationships. *Ann Bot*. 1997;79(3):251–257.
89. Peng C, Zhang L, Liu J. Developing and validating nonlinear height-diameter models for major tree species of Ontario's boreal forests. *North J Appl For*. 2001;18(3):87–94.
90. Colbert JJ, Schuckers M, Fekedulegn D, Rentch J, MacSiúrtáin M, Gottschalk K. Individual tree basal-area growth parameter estimates for four models. *Ecol Model*. 2004;174(1–2):115–126.

91. Khamis A, Ismail Z, Haron K, Mohammed AT. Nonlinear growth models for modeling oil palm yield growth. *J Mathematics and Statistics*. 2005;1(3):225–233.
92. Hau B, Kosman E. Analytical and theoretical plant pathology comparative analysis of flexible two-parameter models of plant disease epidemics. *Phytopathology*. 2007;97(10):1231–1244.
93. Donoso P, Donoso C, Navarro C, Escobar B. 2006. *Nothofagus dombeyi*. In Donoso C (ed.) Las especies arbóreas de los bosques templados de Chile y Argentina. Autoecología. Marisa Cuneo Ediciones, Valdivia, Chile, p. 423–432.
94. • Lappi J, Bailey RL. A height prediction model with random stand and tree parameters: an alternative to traditional site index methods. *Forest Sci*. 1988;34:907–927. **For the first time, mixed-effects models are applied to forest growth modelling.**
95. Lindstrom MJ, Bates DM. Nonlinear mixed effects models for repeated measures data. *Biometrics*. 1990;46(3):673–687.
96. Gregoire TG, Schabenberger O. Nonlinear mixed-effects modeling of cumulative bole volume with spatially correlated within-tree data. *J Agric Biol Environ Stat*. 1996;1:107–119.
97. Pinheiro JC, Bates DM. Mixed-effects models in S and Splus. Springer-verlag, New York, USA, pp 528. 2000.
98. Pinheiro J, Bates D, DebRoy S, Sarkar D, R Core Team. nlme: linear and nonlinear mixed effects models. <https://CRAN.R-project.org/package=nlme>. R package version 3.1-152. 2021.
99. R Core Team. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org>. 2020.
100. Salas C, Ene L, Gregoire TG, Næsset E, Gobakken T. Modelling tree diameter from airborne laser scanning derived variables: A, comparison of spatial statistical models. *Remote Sensing of Environment*. 2010;114(6):1277–1285.
101. Wang M, Rennols K. Tree diameter distribution modelling: introducing the logit-logistic distribution. *Can J For Res*. 2005;35:1305–1313.
102. Jolicoeur P. Human stature: which growth model? *Growth, Development & Aging*. 1991;55:129–132.
103. Huang S, Titus SJ, Wiens DP. Comparison of nonlinear height-diameter functions for major Alberta tree species. *Can J For Res*. 1992;22(9):1297–1304.
104. Kobe RK. Light gradient partitioning among tropical tree species through differential seedling mortality and growth. *Ecol*. 1999;80(1):187–201.
105. Kobe RK. Sapling growth as a function of light and landscape-level variation in soil water and foliar nitrogen in northern michigan. *Oecologia*. 2006;147(1):119–133.
106. Holste EK, Kobe RK, Vriesendorp CF. Seedling growth responses to soil resources in the understory of a wet tropical forest. *Ecol*. 2011;92(9):1828–1838.
107. Tjørve E, Tjørve KMC. A unified approach to the Richards-model family for use in growth analyses: Why we need only two model forms. *Journal of Theoretical Biology*. 2010;267(3):417–425.
108. Causton DR. A computer program for fitting the Richards function. *Biometrics*. 1969;25(2):401–409.
109. Fresco LFM. A model for plant-growth – estimation of the parameters of the logistic function. *Acta Botanica Neerlandica*. 1973;22:486–489.
110. von Gadow K, Hui G. Modelling Forest Development. Cuvillier Verlag, Göttingen, Germany, pp 211. 1999.
111. Weiskittel AR, Hann DW, Kershaw JA Jr, Vanclay JK. Forest growth and yield modeling. New York: Wiley; 2011, p. 430.
112. Stage AR. How forest models are connected to reality: evaluation criteria for their use in decision support. *Can J For Res*. 2003;33(3):410–421.
113. Soto DP, Jacobs DF, Salas C, Donoso PJ, Fuentes C, Puettmann KJ. Light and nitrogen interact to influence regeneration in old-growth *Nothofagus*-dominated forests in south-central Chile. *For Ecol Manage*. 2017;384:303–313.
114. Pacala SW, Canham CD, Saponara J, Silander JA, Kobe RK, Ribbens E. Forest models defined by field measurements: estimation, error analysis and dynamics. *Ecol Appl*. 1996;6(1):1–43.
115. Coates KD, Burton PJ. Growth of planted tree seedlings in response to ambient light levels in northwestern interior cedar-hemlock forests of British Columbia. *Can J For Res*. 1999;29:1347–1382.
116. Soto DP, Donoso PJ, Salas C, Puettmann KJ. Light availability and soil compaction influence the growth of underplanted *Nothofagus* following partial shelterwood harvest and soil scarification. *Can J For Res*. 2015;45:998–1005.
117. Twilley RR, Rivera-Monroy VH, Chen R, Botero L. Adapting an ecological mangrove model to simulate trajectories in restoration ecology. *Mar Pollut Bull*. 1999;37(8):404–419.
118. Petit B, Montagnini F. Growth equations and rotation ages of ten native tree species in mixed and pure plantations in the humid neotropics. *For Ecol Manage*. 2004;2004(2):243–257.
119. Vaverková MD, Radziemska M, Bartoň S, Cerdà A, Koda E. The use of vegetation as a natural strategy for landfill restoration. *Land Degradation & Development*. 2018;29(10):3674–3680.
120. Niklas K. Plant allometry: the scaling of form and process. University of Chicago Press, USA, pp 395. 1994.
121. Enquist B, Brown JH, West GB. Allometric scaling of plant energetics and population density. *Nature*. 1998;395:163–165.
122. West GB, Enquist BJ, Brown JH. A general model for the structure and allometry of plant vascular systems. *Nature*. 1999;400:664–667.
123. Curtis RO. Height-diameter and height-diameter-age equations for second-growth Douglas-fir. *Forest Sci*. 1967;13(4):365–375.
124. Mehtätalo L, de Miguel S, Gregoire TG. Modeling height-diameter curves for prediction. *Can J For Res*. 2015;45:826–837.
125. Mäkelä A. Derivation of stem taper from the pipe theory in a carbon balance framework. *Tree Physiol*. 2002;22(13):891–905.
126. García O. Dynamic modelling of tree form. *Math.Comput For Nat-Res Sci*. 2015;7:9.
127. Rees M, Osborne CP, Woodward FI, Hulme SP, Turnbull LA, Taylor SH. Partitioning the components of relative growth rate: how important is plant size variation? *The American Naturalist*. 2010;176(6):152–161.
128. Pommerening A, Muszta A. Methods of modelling relative growth rate. *Forest Ecosystems*. 2015;2:5.
129. • Elliston AM, Dennis B. Paths to statistical fluency for ecologists. *Frontiers in Ecology and the Environment*. 1996;8(7):362–370. **Addresses the importance of mathematical and statistical knowledge in ecology.**
130. García O. Problemas y modelos en el manejo de las plantaciones forestales. Tesis Ingeniero Forestal, Universidad de Chile, Santiago, Chile, pp 64. 1968.
131. Leary RA, Skog KE. A computational strategy for system identification in ecology. *Ecology*. 1972;53(5):969–973.
132. Leary RA. Design. In a generalized forest growth projection system applied to the Lake states region. USDA For. Serv., Gen. Tech. Rep. NC-49. Sant Paul, Minnesota, USA, pp 5–15. 1979.

133. ●● Hamlin D, Leary RA. Methods for using an integro-differential equation as a model of tree height growth. *Can J For Res.* 1987;17(5):353–356. **The study gives mathematical and statistical details for fitting differential equations using tree growth data.**
134. Dennis B. Statistics and the scientific method in ecology. The nature of scientific evidence: statistical, philosophical, and

empirical considerations. The University of Chicago Press, Chicago, IL, USA, pp 327–378. In: Taper ML and Lele SR, editors; 1996.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Affiliations

Christian Salas-Eljatib^{1,2,3}  · Lauri Mehtätalo⁴ · Timothy G. Gregoire⁵ · Daniel P. Soto⁶  · Rodrigo Vargas-Gaete^{7,8}

¹ Centro de Modelación y Monitoreo de Ecosistemas, Universidad Mayor, Santiago, Chile

² Vicerrectoría de Investigación y Postgrado, Universidad de La Frontera, Temuco, Chile

³ Departamento de Silvicultura y Conservación de la Naturaleza, Universidad de Chile, Santiago, Chile

⁴ Natural Resources Institute Finland (Luke), Joensuu, Finland

⁵ School of the Environment, Yale University, New Haven, CT, USA

⁶ Departamento de Ciencias Naturales y Tecnología, Universidad de Aysén, Coyhaique, Chile

⁷ Laboratorio de Biometría, Departamento de Ciencias Forestales, Universidad de La Frontera, Temuco, Chile

⁸ Centro Nacional de Excelencia para la Industria de la Madera (CENAMAD), Pontificia Universidad Católica de Chile, Santiago, Chile